

Rumex acetosella L.. (Polygonaceae)
Sheep Sorrel, Red Sorrel

Description. Dioecious or polygamous perennials, colonial, from slender rhizomes. Stems 1 to 5 dm tall, erect, simple, glabrous. Leaves alternate, the stipules modified into prominent sheaths, lower leaves petiolate, 2 to 6 cm long, 1 to 2.5 cm wide, blade shape variable, linear to ovate, hastate to sagittate, the terminal lobe much longer than the basal lobes, acute to obtuse, margins entire, the cauline leaves reduced upwards, short-petiolate to sessile, the blades with 1 or no lobes, upper ones sheath-like. Inflorescence 5 to 15 cm long, an open panicle, the branches slender, ascending, glabrous, yellowish to reddish. Flowers unisexual, in clusters, sessile to pedicellate, often somewhat pendulous, perianth parts in two whorls of 3 each, 0.5 to 2 mm long, greenish to reddish, fused near the base, the inner ones expanding and somewhat enclosing the achene in fruit; staminate flowers with 6 stamens; pistillate flowers with 1 superior ovary, the styles 3, the ovary 3-angled, with 1 locule. Fruit a 3-angled achene, 1-2.5 mm long, yellowish to brownish red. In California, flowering from March to July. (Gleason and Cronquist 1991, Hickman 1993, Kaul 1986, Munz 1959, Welsh 1984).

Geographic distribution. A native of Eurasia, sheep sorrel has become naturalized throughout North America, South America, Australia, New Zealand, Japan, southern Africa and Hawaii (Clapham et al. 1982, Fernald 1950, Gleason and Cronquist 1991, Gobbi 1995, Kaul 1986, Munz 1959, Ohwi 1965, Wagner et al. 1990, Webb et al. 1988).

Sheep sorrel was first reported from California (without locality) by Watson (1880), but probably was widely distributed prior to the late 1800s (Jepson 1914, Robbins 1940). It has been reported only from Santa Cruz and Santa Rosa islands (Junk et al. 1997) but occurs widely throughout most mainland California counties (Anonymous 1998, Hickman 1993).

Reproductive and vegetative biology. Sheep sorrel is dioecious, pollinated by wind, but occasionally visited (for pollen) by insects (Proctor et al. 1996, Richards 1986). Seeds may remain dormant for up to 20 years (Tsuyuzaki and Kanda 1996).

Although sheep sorrel represents a polyploid complex (2x, 4x, 6x, and 8x) worldwide, chromosome number does not appear related to ecological distribution nor does it have an effect on sex ratios (Love 1944, Richards 1986). However, because of clonal growth, sex ratios in some populations are often skewed toward one of the two sexes (Houssard et al. 1994, Houssard and Escarre 1995, Richards 1986). Staminate plants are generally shorter, and flower and senesce early, relative to pistillate plants. Pistillate plants are usually larger and have relatively higher reproductive effort, based on flower number. However, Putwain and Harper (1972) showed that the rarer sex often had higher productivity and reproductive capacity, measured either as pollen or seed output. In most cases, ramet growth from root buds contributed much more to population expansion than annual seed output and recruitment.

Initial establishment of *Rumex acetosella* by seed is relatively rare, but appears favored by reduced competition (at least from grasses) and relatively low nutrient levels (Harper 1977, Mamolos et al. 1995, Putwain et al. 1968). Sheep sorrel is a non-mycorrhizal species, and can expand in the absence of competitive mycorrhizal species after fungicide treatment on nutrient-poor soils (Newsham et al. 1995). Ecotypic variation, combined with phenotypic plasticity, can

result in different genotypes dominating stages of old-field succession, resulting in long-term persistence as a species (Escarre et al. 1994, Houssard and Escarre 1995, Korpelainen 1993).

Ecological distribution. Sheep sorrel is generally found on moist soils of waste places, open, disturbed sites, or vernal moist meadows and grasslands (Fernald 1950, Gleason and Cronquist 1991, Kaul 1986, Munz 1959, Welsh 1984). *Rumex acetosella* has also been reported from coastal prairie and coastal scrub communities (Heady et al. 1988). In South America, it has been reported as a common invasive after grassland fires (Gobbi 1995).

Weed status. Sheep sorrel is not considered a noxious weed in agricultural or horticultural practice, at least at a global level (not listed by Holm et al. 1977), nor is it considered a noxious weed by the State Dept. of Food and Agriculture (Anonymous 1996). However, it is listed for the United States in Lorenzi and Jeffery (1987) and is considered a noxious weed in Australia (Lemerle 1996).

Microbial pathogens. No literature was found that reported *Rumex acetosella* as a host to fungal pathogens. It has been reported as a host to the tomato spotted leaf virus (Bitterlich and McDonald 1993). Migliore et al (1997) documented the uptake of sulphadimethoxine, an antimicrobial compound used in the livestock industry, in sheep sorrel and the potential for ingestion by cattle.

Insect pathogens. No literature was found that reported insects detrimental to *Rumex acetosella*.

Herbicide control. Harper (1977) reported that sheep sorrel in experimental plots was resistant to paraquat, but effectively controlled by 2,4-D and Tordon. No other literature was found that specifically addressed herbicide control of *Rumex acetosella*. Lorenzi and Jeffery (1987), however, recommended mechanical removal or spot treatment with 2,4-D plus dicamba for low density infestations and glyphosate for large infested areas.

Literature Cited

- Anonymous. 1996. Exotic pest plants of greatest ecological concern in California as of August 1996. California Exotic Pest Plant Council. 8 pp.
- Anonymous. 1998. USDA Plants Database, Baton Rouge, LA. URL = usda.plants.gov.
- Arnold, T. and B. de Wet. 1993. Memoir 62. Plants of southern Africa: names and distribution. National Botanical Institute, Pretoria. 825 pp.
- Bitterlich, I. and L. McDonald. 1993. The prevalence of tomato spotted wilt virus in weeds and crops in southwestern British Columbia. Canadian Plant Disease Survey. 73: 137-142.
- Clapham, A., T. Tutin, and E. Warburg. 1962. Flora of the British Isles. Cambridge University Press, Cambridge. 1269 pp.
- Escarre, J., C. Houssard, and J. Thompson. 1994. An experimental study of the role of seedling density and neighbor relatedness in the persistence of *Rumex acetosella* in an old-field succession. Canadian Journal of Botany. 72: 1273-1281.

- Fernald, M. 1950. Gray's manual of botany. 8th edition. American Book Company, New York. 1632 pp.
- Gleason, H. and A. Cronquist. 1991. Manual of the vascular plants of northeastern United States and Adjacent Canada. 2nd edition. New York Botanic Garden, Bronx. 910 pp.
- Gobbi, M. 1995. Post-fire recovery and invasion by alien plant species in a South American woodland-steppe ecotone. pp. 105-115. In Pysek, P., (Ed.). Plant invasions: General aspects and special problems. Academic Publishing, The Hague, Netherlands. 263 pp.
- Harper, J. 1977. Population biology of plants. Academic Press, London. 892 pp.
- Heady, H., T. Foin, M. Hektner, D. Taylor, M. Barbour, and W. Barry. 1988. Coastal prairie and northern coastal scrub. pp. 733-760. In Barbour, M. and J. Major (eds.). Terrestrial vegetation of California. California Native Plant Society, Sacramento. 1002 pp.
- Hickman, J. 1993. Polygonaceae. pp. 854-895. In Hickman, J. (ed.). The Jepson manual: Vascular plants of California. University of California Press, Berkeley. 1400 pp.
- Holm, L., D. Plucknett, J. Pancho, and J. Herberger. 1977. The world's worst weeds: distribution and ecology. University Press of Hawaii, Honolulu. 609 pp.
- Houssard, C. and J. Escarre. 1995. Variation and covariation among life-history traits in *Rumex acetosella* from a successional old-field gradient. *Oecologia*. 102: 70-80.
- Houssard, C., J. Thompson, and J. Escarre 1994. Do sex-related differences in response to environmental variation influence the sex-ratio in the dioecious *Rumex acetosella*? *Oikos*. 70: 80-90.
- Jepson, W. 1914. A flora of California. Volume 1. Part 4. pp. 369-464. Jepson Herbarium, University of California, Berkeley.
- Junak, S., S. Chaney, R. Philbrick, and R. Clark. 1997. A checklist of vascular plants of Channel Islands National Park. Southwest Parks and Monuments Association, Tucson, AZ. 43 pp.
- Kaul, R. Polygonaceae. pp. 213-245. In Great Plains Flora Association. 1986. Flora of the Great Plains. University of Kansas, Lawrence. 1392 pp.
- Korpelainen, H. 1993. Vegetative growth in *Rumex acetosella* (Polygonaceae) originating from different geographic regions. *Plant Systematics and Evolution*. 188: 115-123.
- Lemerle, D., T. Yuan, G. Murray, and S. Morris. 1996. Survey of weeds and diseases in cereal crops in the southern wheat belt of New South Wales. *Australian Journal of Experimental Agriculture*. 36: 545-554.
- Lorenzi, H. and L. Jeffery. 1987. Weeds of the United States and their control. Van Nostrand Company, New York. 355 pp.
- Love, A. 1944. Cytogenetic studies on *Rumex* subgenus *Acetosella*. *Hereditas* 30: 1-136.
- Mamolos, A., D. Veresoglou, and N. Barbayiannis. 1995. Plant species abundance and tissue concentrations of limiting nutrients in low-nutrient grasslands: A test of competition theory. *Journal of Ecology*. 83: 485-495.
- Migliore, L., C. Civitareale, G. Brambilla, S. Cozzolino, P. Casoria, and L. Gaudio. 1997. Effects of sulphadimethoxine on cosmopolitan weeds (*Amaranthus retroflexus* L., *Plantago major* L. and *Rumex acetosella* L.). *Agriculture Ecosystems & Environment*. 65: 163-168.

- Munz, P. 1959. A flora of California. University of California Press, Berkeley. 1681 pp.
- Newsham, K., A. Watkinson, H. West, and A. Fitter. 1995. Symbiotic fungi determine plant community structure: Changes in a lichen-rich community induced by fungicide application. *Functional Ecology* 9: 442-447.
- Ohwi, J. 1965. Flora of Japan. Smithsonian Institution, Washington D.C. 1066 pp.
- Proctor, M. P. Yeo, and A. Lack. 1996. The natural history of pollination. Timber Press, Portland, Oregon. 479 pp.
- Putwain, P., D. Machin, and J. Harper. 1968. Studies in the dynamics of plant populations. II. Components and regulation of a natural population of *Rumex acetosella* L. *Journal of Ecology* 56: 421-431.
- Putwain, P. and J. Harper. 1972. Studies in the dynamics of plant populations. V. Mechanisms governing the sex ratio in *Rumex acetosa* and *R. acetosella*. *Journal of Ecology* 60: 113-129.
- Richards, A. 1986. Plant breeding systems. Unwin Hyman, London. 529 pp.
- Robbins, W. 1940. Alien plants growing without cultivation in California. Agricultural Experiment Station. Bulletin 637. University of California, Berkeley. 128 pp.
- Tsuyuzaki, S. and F. Kanda. 1996. Revegetation patterns and seedbank structure on abandoned pastures in northern Japan. *American Journal of Botany* 83: 1422-1428.
- Wagner, W., D. Herbst, and S. Sohmer. 1990. Manual of the flowering plants of Hawaii. 1853 pp.
- Watson, S. 1880. Geological Survey of California. Botany Volume 2. John Wilson, University Press, Cambridge, Massachusetts. 559 pp.
- Webb, C., W. Sykes, and P. Garnock-Jones. 1988. Flora of New Zealand. Volume 4. Naturalized pteridophytes, gymnosperms, dicotyledons. Department of Scientific and Industrial Research, Christchurch. 1365 pp.
- Welsh, S. 1984. Utah flora. Polygonaceae. *Great Basin Naturalist* 44: 519-557.